

stilbene dibromide

Stilbene Dibromide: An In-Depth Overview

Stilbene dibromide is a significant organic compound widely studied in the fields of chemistry and molecular synthesis. Recognized for its unique structure and versatile applications, stilbene dibromide serves as a key intermediate in various chemical reactions. This article offers a comprehensive overview of stilbene dibromide, including its chemical properties, synthesis methods, applications, safety considerations, and analytical techniques.

Understanding Stilbene Dibromide

Stilbene dibromide is a dihalogenated derivative of stilbene, a hydrocarbon composed of two benzene rings connected by an ethene bridge. Its chemical formula is typically represented as $C_{14}H_{12}Br_2$, reflecting the addition of bromine atoms. The compound exists in different isomeric forms, primarily the *cis*- and *trans*- configurations, which influence its physical and chemical properties.

Chemical Structure and Properties

Stilbene dibromide features two bromine atoms attached across the double bond of the stilbene backbone. Its key properties include:

- **Physical State:** Usually a crystalline solid or oily liquid depending on purity and temperature.
- **Molecular Weight:** Approximately 370.06 g/mol.
- **Appearance:** Typically colorless to pale yellow crystalline or oily substance.
- **Reactivity:** Highly reactive towards nucleophiles and participates readily in addition reactions.
- **Stability:** Sensitive to light and temperature; storage in dark, cool conditions recommended.

The presence of bromine atoms significantly enhances the compound's reactivity, making it a valuable intermediate in synthesis.

Synthesis of Stilbene Dibromide

The synthesis of stilbene dibromide involves halogenation of stilbene or related compounds. Several methods have been developed to produce high-purity samples suitable for research and industrial applications.

Common Synthesis Routes

1. Direct Bromination of Stilbene:

- React stilbene with bromine (Br_2) in an inert solvent such as carbon tetrachloride or dichloromethane.
- Control reaction conditions to favor dibromination and prevent polybromination.
- Use of a catalyst or light to initiate the reaction, often under controlled temperature.

2. Halogenation Using N-Bromosuccinimide (NBS):

- Employ NBS as a brominating agent in the presence of a radical initiator like AIBN.
- Provides milder reaction conditions and better control over bromination levels.

3. Oxidative Bromination:

- Involves oxidation of bromide ions in situ to bromine, which then reacts with stilbene.
- Typically performed in aqueous media with appropriate oxidants.

Purification Techniques

Post-synthesis, stilbene dibromide often requires purification to attain desired purity levels:

- **Recrystallization:** Using solvents like ethanol or acetic acid to isolate pure crystals.
- **Column Chromatography:** For separating isomers or removing impurities.
- **Distillation:** In cases where the compound is in liquid form, to refine its purity.

Applications of Stilbene Dibromide

The utility of stilbene dibromide spans various domains, primarily due to its reactive double bond and bromine substituents. Its applications include:

1. Organic Synthesis Intermediary

Stilbene dibromide serves as a precursor in synthesizing complex organic molecules. Its brominated double bond allows for subsequent reactions such as:

- Elimination reactions to generate stilbene derivatives with specific properties.
- Formation of dyes, polymers, and pharmaceuticals through nucleophilic substitution.
- Preparation of trans- and cis-isomers for stereochemical studies.

2. Photochemical Studies

Due to its photo-reactive nature, stilbene dibromide is used in studies involving:

- Photoisomerization processes.
- Photophysics and photochemistry to understand light-induced reactions.
- Development of photoresponsive materials.

3. Material Science and Dyes

Its conjugated system and ability to undergo halogenation reactions make it suitable for:

- Designing organic dyes and pigments.
- Developing photochromic and thermochromic materials.

4. Research and Development

In academic research, stilbene dibromide is employed to:

- Investigate reaction mechanisms involving halogenated alkenes.
- Study stereochemical properties of trans- and cis-isomers.
- Explore new synthetic pathways for organic compounds.

Safety and Handling

Handling stilbene dibromide necessitates strict safety precautions due to its chemical reactivity and potential health hazards.

Safety Precautions

- **Personal Protective Equipment:** Use gloves, lab coat, and eye protection.
- **Ventilation:** Conduct reactions in a well-ventilated fume hood.
- **Storage:** Store in a cool, dry, and dark place, in tightly sealed containers.
- **Disposal:** Dispose of waste according to local environmental regulations, avoiding release into the environment.

Potential Hazards

- Bromine compounds are toxic and may cause skin and eye irritation.
- Inhalation of vapors may lead to respiratory issues.
- Flammable solvents used during synthesis pose fire risks.

Analytical Techniques for Characterization

Accurate characterization of stilbene dibromide is vital for confirming purity and structure. Common analytical methods include:

1. Nuclear Magnetic Resonance (NMR) Spectroscopy

- ^1H NMR: Identifies hydrogen environments and confirms double-bond configurations.
- ^{13}C NMR: Provides insights into carbon skeleton and brominated carbons.

2. Infrared (IR) Spectroscopy

- Detects characteristic absorption bands such as C=C stretching and C-Br vibrations.

3. Mass Spectrometry (MS)

- Confirms molecular weight and isotopic patterns due to bromine isotopes.

4. UV-Vis Spectroscopy

- Assesses conjugation and electronic transitions relevant to photochemical applications.

5. Chromatography Techniques

- Thin-layer chromatography (TLC) and high-performance liquid chromatography

(HPLC) for purity assessment.

Conclusion

Stilbene dibromide is a versatile and valuable compound in organic chemistry, offering numerous pathways for synthesis and applications in materials science, photochemistry, and research. Its reactive nature, combined with well-established synthesis and purification methods, makes it an essential intermediate for chemists seeking to develop novel compounds and materials. Proper handling and analytical characterization ensure safe and effective utilization of this compound, paving the way for innovative developments in various scientific disciplines.

References

- Organic Chemistry textbooks and synthesis manuals.
- Scientific journals on halogenated hydrocarbons and photochemical compounds.
- Material safety data sheets (MSDS) for bromine compounds.
- Analytical chemistry handbooks for spectroscopy techniques.

Frequently Asked Questions

What is stilbene dibromide and its primary use in organic chemistry?

Stilbene dibromide is an organic compound derived from stilbene, where two bromine atoms are added across the double bond. It is primarily used as an intermediate in organic synthesis, especially in the preparation of stilbene derivatives and in stereochemical studies.

How is stilbene dibromide synthesized in the laboratory?

Stilbene dibromide is typically synthesized by adding bromine (Br_2) to stilbene in a solvent like carbon tetrachloride or chloroform, resulting in the addition across the double bond and forming the dibromide compound.

What are the key safety considerations when handling stilbene dibromide?

Handling stilbene dibromide requires safety precautions such as working in a well-ventilated area, wearing gloves and eye protection, and avoiding inhalation or skin contact, as it is a brominated organic compound that may be toxic or irritant.

What are the spectroscopic features used to identify stilbene dibromide?

Stilbene dibromide can be characterized by its distinctive IR absorption bands (notably C-Br stretches), NMR signals showing the addition of bromines across the double bond, and mass spectrometry confirming its molecular weight and fragmentation pattern.

How does the stereochemistry of stilbene dibromide influence its chemical properties?

The stereochemistry (cis or trans isomers) of stilbene dibromide affects its physical properties, reactivity, and biological activity. The trans isomer is generally more stable and less reactive, influencing how it participates in subsequent reactions.

What are recent research trends involving stilbene dibromide?

Recent research trends include exploring its use in synthesizing novel organic materials, studying its stereochemical behavior in reactions, and investigating its potential as a precursor in developing pharmaceuticals and functional materials.

Additional Resources

Stilbene Dibromide: A Comprehensive Guide to Its Structure, Synthesis, and Applications

Stilbene dibromide is an intriguing organic compound that captures the attention of chemists and researchers due to its unique structure and versatile applications. As a derivative of stilbene, a well-known aromatic alkene, stilbene dibromide features two bromine atoms attached to the stilbene backbone, imparting distinctive chemical properties that make it valuable in various fields, from organic synthesis to material science. This article aims to provide a detailed exploration of stilbene dibromide, including its chemical structure, synthesis methods, characterization techniques, and practical uses.

Understanding the Basics of Stilbene Dibromide

What Is Stilbene Dibromide?

At its core, stilbene dibromide is a halogenated alkene derived from stilbene, a molecule characterized by two phenyl groups attached to an ethene chain. The "dibromide" suffix indicates that two bromine atoms are added

across the double bond, converting it into a vicinal dibromide.

Chemical Formula: $C_{14}H_{14}Br_2$

Molecular Weight: Approximately 341.03 g/mol

Structural Formula:

```

Ph-CH=CH-Ph (stilbene backbone)

|

Br

|

Br

```

(Note: The bromines are attached to the carbons of the former double bond, converting it into a dibromide.)

Structural Features and Isomerism

Stilbene dibromide exists in different isomeric forms, primarily cis and trans configurations:

- Cis-Isomer: Both bromine atoms are on the same side of the double bond.
- Trans-Isomer: Bromine atoms are on opposite sides.

The isomerism can influence the compound's physical and chemical properties, including melting points, reactivity, and optical activity. Typically, the trans-isomer is more thermodynamically stable due to minimized steric hindrance.

Synthesis of Stilbene Dibromide

General Methods

The synthesis of stilbene dibromide primarily involves the bromination of stilbene through electrophilic addition reactions. This process is generally straightforward and can be conducted under controlled laboratory conditions.

Common Synthetic Route

1. Starting Material: Stilbene (either synthetic or extracted from natural sources)
2. Reagent: Bromine (Br_2), often in an organic solvent such as carbon tetrachloride (CCl_4) or dichloromethane (DCM)
3. Reaction Conditions: Room temperature to mild heating, with stirring

Reaction Steps:

- Step 1: Dissolve stilbene in an inert solvent.
- Step 2: Slowly add bromine solution while maintaining constant stirring.
- Step 3: Bromine adds across the double bond in an electrophilic addition manner, forming the dibromide.
- Step 4: The reaction is monitored until the color of bromine disappears, indicating completion.
- Step 5: The product is isolated via filtration, washing, and recrystallization.

Considerations and Variations

- Control of stereochemistry: Reaction conditions can influence the cis/trans ratio.
- Use of catalysts or additives: In some cases, catalysts like FeBr_3 can be employed to control reaction rate.
- Purification: Techniques such as column chromatography or recrystallization are used to purify the dibromide.

Characterization Techniques

Accurate identification and analysis of stilbene dibromide involve various spectroscopic and analytical methods:

1. NMR Spectroscopy

- ^1H NMR: Provides information about the hydrogen environment, confirming the addition of bromines and the stereochemistry.
- ^{13}C NMR: Offers insights into carbon bonding environments, confirming the substitution pattern.

2. Infrared (IR) Spectroscopy

- Identifies characteristic vibrational bands corresponding to C-Br bonds and aromatic rings.

3. Mass Spectrometry (MS)

- Confirms molecular weight and the presence of bromine isotopic patterns.

4. Melting Point and Optical Rotation

- Physical properties help differentiate between isomers and assess purity.

Applications of Stilbene Dibromide

While stilbene dibromide may not be as widely known as other halogenated derivatives, it plays a significant role in various scientific and industrial

contexts.

1. Organic Synthesis Intermediate

- Building Block: Stilbene dibromide serves as an intermediate in synthesizing more complex organic molecules, including dyes, pharmaceuticals, and polymers.
- Cross-coupling Reactions: The dibromide can undergo coupling reactions such as Suzuki or Heck couplings, facilitating the construction of extended aromatic systems.

2. Photoresponsive Materials

- Photoisomerization Studies: Due to its conjugated structure, stilbene dibromide can participate in studies related to light-induced isomerization, useful in developing photoresponsive materials.

3. Dyes and Pigments

- As a derivative of stilbene, which forms the basis of certain dyes, dibromides can modify optical properties, leading to potential applications in dye synthesis or as fluorescent probes.

4. Research and Educational Use

- Model Compound: Used in academic settings to demonstrate electrophilic addition, stereoisomerism, and halogenation reactions.

Safety and Handling

Handling stilbene dibromide requires adherence to safety protocols:

- Toxicity: Brominated organic compounds can be toxic; avoid inhalation or skin contact.
- Protective Equipment: Use gloves, goggles, and lab coats.
- Ventilation: Conduct reactions in a fume hood.
- Disposal: Follow proper chemical waste disposal regulations to prevent environmental contamination.

Future Perspectives and Research Directions

Research into stilbene dibromide continues to evolve, with current focus areas including:

- Development of Novel Materials: Incorporating dibromide derivatives into organic electronics and photoresponsive systems.
- Green Chemistry Approaches: Exploring more sustainable synthesis methods

with less hazardous reagents.

- **Biological Activity:** Investigating potential bioactivity or pharmacological properties, given the biological relevance of stilbene derivatives.

Conclusion

Stilbene dibromide stands out as a versatile and important compound in organic chemistry, bridging the gap between fundamental reactions and practical applications. Its distinctive structure, characterized by the addition of bromines across the stilbene double bond, makes it a valuable intermediate for synthesis, material development, and scientific research. Understanding its synthesis, properties, and potential uses not only enriches the knowledge base but also opens avenues for innovation in chemical sciences.

Whether you're a seasoned chemist or a student exploring halogenated compounds, stilbene dibromide offers a compelling example of how molecular modifications can influence properties and enable diverse applications. As research advances, expect to see even more innovative uses for this fascinating compound.

Stilbene Dibromide

stilbene dibromide: Microscale Organic Laboratory Dana W. Mayo, Ronald M. Pike, David C. Forbes, 2010-01-12 This is a laboratory text for the mainstream organic chemistry course taught at both two and four year schools, featuring both microscale experiments and options for scaling up appropriate experiments for use in the macroscale lab. It provides complete coverage of organic laboratory experiments and techniques with a strong emphasis on modern laboratory instrumentation, a sharp focus on safety in the lab, excellent pre- and post-lab exercises, and multi-step experiments. Notable enhancements to this new edition include inquiry-driven experimentation, validation of the purification process, and the implementation of greener processes (including microwave use) to perform traditional experimentation.

stilbene dibromide: Advanced Organic Chemistry Francis A. Carey, Richard J. Sundberg, 2007-06-13 The two-part, fifth edition of Advanced Organic Chemistry has been substantially revised and reorganized for greater clarity. The material has been updated to reflect advances in the field since the previous edition, especially in computational chemistry. Part A covers fundamental structural topics and basic mechanistic types. It can stand-alone; together, with Part B: Reaction and Synthesis, the two volumes provide a comprehensive foundation for the study in organic chemistry. Companion websites provide digital models for study of structure, reaction and selectivity for students and exercise solutions for instructors.

stilbene dibromide: Organic Chemistry; Or, Chemistry of the Carbon Compounds Victor von Richter, 1922

stilbene dibromide: Stereochemistry Mr. Rohit Manglik, 2024-03-10 EduGorilla Publication is a trusted name in the education sector, committed to empowering learners with high-quality study materials and resources. Specializing in competitive exams and academic support, EduGorilla provides comprehensive and well-structured content tailored to meet the needs of students across

various streams and levels.

stilbene dibromide: Organic Chemistry: Chemistry of the carbocyclic compounds; tr. from the 11th German ed. by E. E. Fournier d'Albe, 1922 Victor von Richter, 1922

stilbene dibromide: Introduction to the Study of Organic Chemistry John Wade, 1898

stilbene dibromide: Organic Reaction Mechanisms 1984 A. C. Knipe, W. E. Watts, 2008-04-30

The only book series to summarize the latest progress on organic reaction mechanisms, *Organic Reaction Mechanisms*, 1984 surveys the development in understanding of the main classes of organic reaction mechanisms reported in the primary scientific literature in 1984. The 20th annual volume in this highly successful series highlights mechanisms of stereo-specific reactions. Reviews are compiled by a team of experienced editors and authors, allowing advanced undergraduates, graduate students, postdocs, and chemists to rely on the volume's continuing quality of selection and presentation.

stilbene dibromide: *Some Modern Methods of Organic Synthesis* W. Carruthers, 1971-10-31

stilbene dibromide: *A Text-book of Organic Chemistry* August Bernthsen, 1889

stilbene dibromide: Journal of the Chemical Society , 1875

stilbene dibromide: Journal of the Chemical Society Chemical Society (Great Britain), 1900

Titles of chemical papers in British and foreign journals included in Quarterly journal, v. 1-12.

stilbene dibromide: *Journal - Chemical Society, London* Chemical Society (Great Britain), 1897

stilbene dibromide: *Oxygen Radicals in Chemistry and Biology* Wolf Bors, Manfred Saran, David Tait, 2011-06-15 No detailed description available for *Oxygen Radicals in Chemistry and Biology*.

stilbene dibromide: Organic Chemistry, Volume Two Frank C. Whitmore, 2011-12-14 This volume discusses the properties of aromatic compounds, alkaloids and other types of heterocyclic compounds, organophosphorus, and organometallic compounds.

stilbene dibromide: *Carbocyclic and heterocyclic series* Victor von Richter, 1900

stilbene dibromide: *Solvents and Solvent Effects in Organic Chemistry* Christian Reichardt, Thomas Welton, 2011-08-04 Now in its 4th edition, this book remains the ultimate reference for all questions regarding solvents and solvent effects in organic chemistry. Retaining its proven concept, there is no other book which covers the subject in so much depth, the handbook is completely updated and contains 15% more content, including new chapters on Solvents and Green chemistry, Classification of Solvents by their Environmental Impact, and Ionic Liquids. An essential part of every organic chemist's library.

stilbene dibromide: *Physico-chemical Constants* Thomas Carnelley, 1885

stilbene dibromide: *Foundations of Organic Chemistry* David R. Dalton, 2011-07-12 This book differs from other organic chemistry textbooks in that it is not focused purely on the needs of students studying premed, but rather for all students studying organic chemistry. It directs the reader to question present assumptions rather than to accept what is told, so the second chapter is largely devoted to spectroscopy (rather than finding it much later on as with most current organic chemistry textbooks). Additionally, after an introduction to spectroscopy, thermodynamics and kinetics, the presentation of structural information of compounds and organic families advances from hydrocarbons to alcohols to aldehydes and ketones and, finally, to carboxylic acids.

stilbene dibromide: *A text-book of organic chemistry, tr. by G. M'Gowan* August Bernthsen, 1894

stilbene dibromide: Victor Von Richter's Organic Chemistry: Carbocyclic and heterocyclic series Victor von Richter, 1900

Related to stilbene dibromide

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula $C_6H_5CH=CHC_6H_5$. Classified as a diarylethene, it features

Stilbene - 103-30-0, C₁₄H₁₂, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C₁₄H₁₂ | CID 11502 - PubChem Stilbene | C₁₄H₁₂ | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C₆-C₂

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula C₆H₅CH=CHC₆H₅. Classified as a diarylethene, it features

Stilbene - 103-30-0, C₁₄H₁₂, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C₁₄H₁₂ | CID 11502 - PubChem Stilbene | C₁₄H₁₂ | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence against There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C₆-C₂

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the

important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula $C_6H_5CH=CHC_6H_5$. Classified as a diarylethene, it features

Stilbene - 103-30-0, C14H12, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C14H12 | CID 11502 - PubChem Stilbene | C14H12 | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence against There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C6-C2

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula $C_6H_5CH=CHC_6H_5$. Classified as a diarylethene, it features

Stilbene - 103-30-0, C14H12, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C14H12 | CID 11502 - PubChem Stilbene | C14H12 | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence against There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for

their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C6-C2

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula $C_6H_5CH=CHC_6H_5$. Classified as a diarylethene, it features

Stilbene - 103-30-0, C14H12, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C14H12 | CID 11502 - PubChem Stilbene | C14H12 | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C6-C2

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

(E)-Stilbene - Wikipedia (E)-Stilbene, commonly known as trans-stilbene, is an organic compound represented by the condensed structural formula $C_6H_5CH=CHC_6H_5$. Classified as a diarylethene, it features

Stilbene - 103-30-0, C14H12, density, melting point, boiling point Stilbene - cas 103-30-0, synthesis, structure, density, melting point, boiling point

Stilbenes, a Versatile Class of Natural Metabolites for Stilbenes are polyphenolic allelochemicals synthesized by plants, especially grapes, peanuts, rhubarb, berries, etc., to defend themselves under stressful conditions. They are now exploited

Stilbene | C14H12 | CID 11502 - PubChem Stilbene | C14H12 | CID 11502 - structure, chemical names, physical and chemical properties, classification, patents, literature, biological activities, safety/hazards/toxicity

(Z)-Stilbene - Wikipedia Stilbene is used in manufacture of dyes and optical brighteners, and also as a phosphor and a scintillator. Stilbene is one of the gain mediums used in dye lasers

Polyphenol Stilbenes: Molecular Mechanisms of Defence There is convincing data that stilbene compounds, a diverse group of natural defence phenolics, abundant in grapes, berries, and conifer bark waste, may confer a protective effect against

Stilbene - an overview | ScienceDirect Topics Natural stilbenes are polyhydroxyphenolic compounds formed through the substitution of hydrogens in different positions of benzene rings by hydroxyl groups with stilbene as the basic

Stilbenes: a journey from folklore to pharmaceutical innovation Stilbenes, polyphenolic compounds synthesized by plants under stress conditions, have garnered significant attention for their therapeutic potential, bridging ancient wisdom with

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. Structurally, they have a C6-C2

Stilbenes: Source plants, chemistry, biosynthesis, pharmacology Stilbenes are some of the important phenolic compounds originating from plant families like Vitaceae, Leguminaceae, Gnetaceae, and Dipterocarpaceae. S

Related Articles

- [acord 25 fillable](#)
- [marine science the dynamic ocean](#)
- [torque chart for metric bolts pdf](#)

[Back to Home](#)