

nursing diagnosis for activity intolerance

nursing diagnosis for activity intolerance is a critical assessment tool used by nurses to identify patients who are experiencing a reduced ability to perform or sustain activities necessary for daily living. Recognizing activity intolerance early allows healthcare professionals to develop targeted interventions aimed at improving patients' functional capacity, promoting independence, and preventing further deterioration. This comprehensive article explores the definition, etiology, signs and symptoms, assessment methods, nursing diagnoses, interventions, and evaluation strategies associated with activity intolerance, providing valuable insights for nursing practice and optimizing patient outcomes.

Understanding Nursing Diagnosis for Activity Intolerance

Definition of Activity Intolerance

Activity intolerance is a clinical condition characterized by an inadequate physiological or psychological response to activity, resulting in fatigue, discomfort, or inability to perform daily tasks. It signifies a diminished capacity to meet the demands of physical or mental exertion, which may be temporary or chronic.

Importance of Nursing Diagnosis for Activity Intolerance

Accurate nursing diagnosis for activity intolerance guides the development of individualized care plans. It ensures that interventions address the root causes, whether they are related to cardiovascular, respiratory, musculoskeletal, or psychological factors. Proper diagnosis also facilitates early rehabilitation, enhances patient safety, and improves quality of life.

Etiology and Contributing Factors

Understanding the causes of activity intolerance is essential in forming effective nursing diagnoses and interventions. These factors may be physiological, psychological, or environmental.

Physiological Causes

- Cardiovascular conditions (e.g., heart failure, ischemic heart disease)
- Respiratory disorders (e.g., chronic obstructive pulmonary disease, asthma)

- Muskuloskeletal limitations (e.g., arthritis, osteoporosis)
- Deconditioning due to prolonged bed rest or inactivity
- Anemia or other hematologic issues
- Metabolic imbalances (e.g., hypothyroidism, diabetes)

Psychological and Emotional Causes

- Anxiety and depression
- Fear of injury or pain during activity
- Lack of motivation or depression

Environmental and Social Factors

- Lack of access to assistive devices or mobility aids
- Unsupportive living environments
- Limited social support or caregiver assistance

Signs and Symptoms of Activity Intolerance

Early recognition of activity intolerance is vital for prompt intervention. Common signs and symptoms include:

- Fatigue after minimal exertion
- Shortness of breath or dyspnea during activity
- Dizziness or lightheadedness
- Palpitations or irregular heartbeat
- Muscle weakness or cramping
- Decreased endurance over time
- Pallor or cyanosis
- Confusion or mental status changes in severe cases

Assessment Strategies for Activity Intolerance

Proper assessment is the cornerstone of establishing an accurate nursing diagnosis. The following strategies are commonly employed:

Subjective Data Collection

- Patient history regarding fatigue, exertion levels, and previous activity tolerance
- Patient reports of dyspnea, chest pain, or discomfort during activity
- Emotional state, motivation, and psychological well-being
- Impact of activity limitations on daily life

Objective Data Collection

- Vital signs at rest and during activity (heart rate, blood pressure, respiratory rate)
- Observation of patient's gait, posture, and mobility
- Measurement of oxygen saturation levels
- Musculoskeletal assessment for weakness or joint limitations
- Functional assessments such as the Six-Minute Walk Test

Tools and Scales

- Borg Rating of Perceived Exertion Scale
- New York Heart Association (NYHA) Functional Classification
- Karnofsky Performance Status Scale
- Activities of Daily Living (ADL) assessments

Formulating Nursing Diagnoses for Activity Intolerance

Using the assessment data, nurses can formulate specific nursing diagnoses based on the NANDA International taxonomy. The most common nursing diagnosis related to activity intolerance is:

- **Activity Intolerance:** Insufficient physiological or psychological energy to meet demands of activity.

Related factors may include:

- Decreased cardiac output
- Impaired gas exchange
- Musculoskeletal impairment
- Fatigue related to anemia
- Anxiety or psychological distress

Defining characteristics include:

- Fatigue
- Shortness of breath

- Dizziness
- Muscle weakness

Goals and Expected Outcomes

Establishing clear goals helps in evaluating the effectiveness of nursing interventions. Typical goals include:

- Patient will demonstrate improved tolerance to activity as evidenced by decreased fatigue and dyspnea.
- Patient will participate in prescribed activity levels without adverse signs.
- Patient will verbalize understanding of activity limitations and management strategies.

Nursing Interventions for Activity Intolerance

Interventions should be individualized based on the patient's specific condition, etiology, and severity. Key interventions include:

Promoting Rest and Energy Conservation

- Encourage periods of rest between activities
- Assist with scheduling activities during optimal energy levels
- Educate on activity pacing and proper body mechanics

Enhancing Physical Activity

- Implement graded exercise programs tailored to patient capacity
- Coordinate with physical therapists for rehabilitation plans
- Encourage gradual increase in activity levels

Monitoring and Managing Physiological Parameters

- Regularly assess vital signs during activity

- Administer oxygen therapy if prescribed
- Manage underlying conditions contributing to activity intolerance

Providing Psychological Support

- Address fears related to activity and health status
- Offer counseling or psychological support when needed
- Encourage patient participation in decision-making

Patient Education

- Teach energy conservation techniques
- Instruct on proper use of assistive devices
- Educate about signs of overexertion and when to seek help
- Discuss importance of adherence to prescribed activity plans

Evaluation of Nursing Outcomes

Effective evaluation measures whether the nursing interventions have achieved the desired outcomes. Indicators include:

- Decreased fatigue and dyspnea during activity
- Increased participation in daily activities
- Patient reports improved energy levels
- Vital signs remain stable during activity
- Patient demonstrates understanding of activity management strategies

Regular re-assessment allows for modification of care plans to ensure ongoing progress and safety.

Conclusion

nursing diagnosis for activity intolerance plays a pivotal role in the holistic care of patients with compromised functional capacity. Recognizing the signs, understanding the underlying causes, and implementing tailored interventions can significantly enhance patients' quality of life. Nurses must perform thorough assessments, develop individualized care plans, and continuously evaluate progress to optimize activity tolerance and promote independence. With a combination of clinical expertise, patient education, and multidisciplinary collaboration, nursing care can effectively address activity intolerance and support patients in achieving their highest possible level of functioning.

Keywords: nursing diagnosis, activity intolerance, assessment, intervention, patient care, fatigue, dyspnea, rehabilitation, energy conservation, functional capacity

Frequently Asked Questions

What are common nursing diagnoses associated with activity intolerance?

Common nursing diagnoses include 'Activity Intolerance' related to decreased oxygenation, fatigue, or pain, often evidenced by verbal reports of exhaustion, decreased activity levels, or abnormal vital signs during activity.

How can nurses assess a patient for activity intolerance?

Nurses can assess activity intolerance through patient history, observing for fatigue, dyspnea, pallor, or dizziness during activity, and measuring vital signs before, during, and after activity to evaluate tolerance levels.

What interventions are effective in managing activity intolerance in nursing care?

Interventions include pacing activities, encouraging rest periods, improving nutrition, providing oxygen therapy if needed, and gradually increasing activity levels to enhance endurance and reduce fatigue.

How does addressing activity intolerance improve patient outcomes?

Managing activity intolerance helps prevent complications like deconditioning, improves functional capacity, enhances quality of life, and promotes independence in daily activities.

What are the expected outcomes when nursing care targets activity intolerance?

Expected outcomes include increased activity tolerance, reduced fatigue and

dyspnea, improved strength and endurance, and the patient's ability to perform ADLs with minimal assistance.

Additional Resources

Nursing diagnosis for activity intolerance is a fundamental component of nursing practice aimed at identifying patients' limitations in performing daily activities due to various physiological, psychological, or environmental factors. Recognizing and accurately diagnosing activity intolerance enables nurses to formulate effective care plans, promote optimal functional status, and improve overall patient outcomes. As a core element of patient-centered care, understanding the intricacies of this nursing diagnosis involves exploring its definition, etiology, signs and symptoms, associated factors, diagnostic criteria, and interventions.

Understanding Activity Intolerance in Nursing Practice

Definition and Conceptual Framework

In nursing, activity intolerance is defined as a state in which an individual experiences insufficient physiological or psychological energy to perform or continue desired daily activities. It is a subjective experience often characterized by fatigue, weakness, or decreased endurance that impairs the ability to carry out routine tasks. The NANDA International (North American Nursing Diagnosis Association) classifies it as a clinical judgment about a person's response to an actual or potential health problem that results in the development of a plan of care aimed at restoring activity levels.

The concept encapsulates both physical and psychological components. Physically, it may manifest as muscle weakness, cardiovascular limitations, or respiratory compromise. Psychologically, factors such as depression, anxiety, or motivation deficits may contribute to or exacerbate activity limitations.

Significance in Patient Care

Addressing activity intolerance is vital because it directly impacts patients' independence, quality of life, and recovery trajectory. For example, post-operative patients or those with chronic illnesses like heart failure or pulmonary diseases often experience activity limitations that can lead to deconditioning, social isolation, and increased risk of complications such as deep vein thrombosis or pneumonia. Early identification and management can prevent deterioration and facilitate rehabilitation.

Etiology and Contributing Factors

Physiological Factors

Several physiological conditions can cause or contribute to activity intolerance, including:

- Cardiovascular impairments: Heart failure, arrhythmias, ischemic heart disease reduce cardiac output, limiting oxygen delivery during exertion.
- Respiratory conditions: Chronic obstructive pulmonary disease (COPD), asthma, pneumonia impair gas exchange, leading to fatigue and dyspnea.
- Musculoskeletal issues: Arthritis, muscular dystrophies, or injuries weaken the musculoskeletal system, decreasing endurance.
- Neurological deficits: Stroke, Parkinson's disease, or multiple sclerosis can impair coordination and stamina.
- Metabolic disturbances: Anemia, diabetes-related complications, or electrolyte imbalances can cause fatigue and reduce activity capacity.

Psychological and Emotional Factors

Mental health status plays a significant role in activity tolerance. Factors include:

- Depression and Anxiety: These can diminish motivation and energy levels.
- Fear of pain or injury: Patients may avoid activity due to fear, leading to deconditioning.
- Cognitive impairments: Cognitive deficits may hinder planning and executing activities.

Environmental and Social Factors

External factors influencing activity levels encompass:

- Lack of social support: Social isolation can diminish activity engagement.
- Environmental barriers: Unsafe or inaccessible environments hinder mobility.
- Cultural beliefs: Cultural attitudes towards activity and rest may affect participation.

Clinical Manifestations and Signs of Activity Intolerance

Subjective Data

Patients may report:

- Persistent fatigue and exhaustion
- Shortness of breath during minimal exertion
- Dizziness or lightheadedness
- Feeling of weakness or heaviness
- Decreased motivation or interest in activities

Objective Data

Nurses observe or measure:

- Decreased endurance during activity tests
- Elevated heart rate (tachycardia) with exertion
- Elevated respiratory rate or oxygen desaturation
- Muscular weakness or tremors
- Post-exertional malaise or prolonged recovery time
- Decreased participation in activities or social withdrawal

Assessment Tools and Measures

Various tools assist in assessing activity tolerance, including:

- Six-Minute Walk Test (6MWT): Measures functional exercise capacity.
- Borg Rating of Perceived Exertion: Subjective scale for fatigue or breathlessness.
- Activity diaries or logs: Track activity levels over time.
- Fatigue scales: Quantify severity and impact on daily life.

Diagnostic Process and Criteria

Formulating the Nursing Diagnosis

NANDA-I provides the diagnostic label "Activity Intolerance" with defining characteristics, related factors, and evidence.

Defining Characteristics include:

- Fatigue
- Dyspnea
- Decreased endurance
- Muscle weakness
- Dizziness
- Post-exertional malaise

Related Factors may involve:

- Cardiac or respiratory impairments
- Anemia
- Musculoskeletal limitations
- Psychological issues such as depression

- Environmental barriers

Evidence or Data can be obtained through patient history, physical examination, and assessment tools.

Examples of Nursing Diagnoses Related to Activity Intolerance

- Risk for activity intolerance (for patients at risk but not yet experiencing symptoms)
- Impaired physical activity tolerance (when manifestations are evident)
- Readiness for enhanced activity tolerance (when patient demonstrates motivation to improve)

Interventions and Management Strategies

Effective management of activity intolerance involves a multifaceted approach tailored to the underlying causes and patient-specific factors.

Primary Nursing Interventions

- Assessment and Monitoring: Regular evaluation of vital signs, activity levels, and fatigue severity.
- Patient Education: Teaching energy conservation techniques, proper pacing, and importance of activity within tolerance.
- Psychosocial Support: Addressing psychological barriers through counseling, motivation, and reassurance.
- Gradual Activity Promotion: Implementing graded exercise programs to enhance endurance safely.
- Environmental Modifications: Ensuring accessible and safe environments to promote mobility.

Collaborative Interventions

- Medical Management: Optimizing treatment for underlying conditions (e.g., medication for heart failure or COPD).
- Physical Therapy: Designing individualized exercise and rehabilitation programs.
- Nutritional Support: Addressing deficiencies (e.g., anemia) that impair activity capacity.
- Psychological Support: Engaging mental health professionals when necessary.

Patient-Centered Goals

- Improving endurance and strength
- Enhancing independence in activities of daily living
- Preventing complications related to inactivity
- Promoting psychological well-being and motivation

Evaluation and Outcomes

Evaluation involves assessing whether interventions have led to:

- Increased activity tolerance
- Reduced fatigue and dyspnea
- Improved functional capacity
- Enhanced quality of life
- Patient ability to perform desired activities independently

Regular reassessment ensures modifications to the care plan as needed and tracks progress toward established goals.

Challenges and Considerations in Nursing Diagnosis of Activity Intolerance

Several challenges can arise in diagnosing activity intolerance accurately:

- Subjectivity of Symptoms: Fatigue and weakness are subjective and vary among individuals.
- Overlap with Other Diagnoses: Symptoms may overlap with other conditions like depression or chronic pain.
- Cultural Factors: Cultural perceptions about activity and rest influence reporting and engagement.
- Patient Variability: Age, comorbidities, and baseline functional status affect assessment and management.

Nurses must employ critical thinking, comprehensive assessment, and holistic approaches to accurately identify and address activity intolerance.

Conclusion

Nursing diagnosis for activity intolerance is a vital aspect of holistic patient care, encompassing a thorough understanding of physiological, psychological, and environmental factors influencing a patient's ability to perform daily activities. Accurate assessment and diagnosis enable targeted interventions, fostering improved endurance, independence, and quality of life. As healthcare continues to evolve toward patient-centered models, nurses' role in recognizing and managing activity intolerance remains indispensable in promoting recovery and well-being across diverse patient

populations.

In summary, addressing activity intolerance requires a comprehensive approach that combines clinical evaluation, patient education, collaborative care, and continuous reassessment. By understanding its underlying causes, manifestations, and management strategies, nurses can significantly impact patient outcomes, ultimately restoring or enhancing their capacity for activity and participation in life.

Nursing Diagnosis For Activity Intolerance

nursing diagnosis for activity intolerance: Nursing Diagnosis Lynda Juall Carpenito-Moyet, 2008 Explains the role of nursing diagnosis in clinical practice; provides information on definitions, characteristics, related factors, and interventions for nursing diagnoses; and offers information on collaborative problems.

nursing diagnosis for activity intolerance: Maternal Newborn Nursing Care Plans Carol Green, 2011-04-15 Maternal Newborn Nursing Care Plans, Second Edition teaches students and professional nurses how to assess, plan, deliver, and evaluate care for normal and abnormal pregnancy, including delivery and recovery for both the mother and baby. This new edition contains more than 65 of the most common and high-risk care plans for maternal and newborn nursing care using the nursing process approach. Organized according to clinical condition, Maternal Newborn Nursing Care Plans, Second Edition provides practical components for each care plan incorporating: * Key nursing activities * Etiologies and risk factors * Signs and symptoms * Diagnostic studies * Medical management * Collaborative problems * Individualized care plans complete with the Nursing Interventions Classification (NIC)

nursing diagnosis for activity intolerance: Application Of Nursing Process and Nursing Diagnosis Marilyn E Doenges, Mary Frances Moorhouse, 2012-12-07 When you understand the whys of each step the nursing process, it's easier easy to understand how to apply them in the real world in which you will practice. Take an interactive, step-by-step approach to developing the diagnostic reasoning and problem-solving skills you need to think like a nurse with the resources you'll find in this unique workbook style text.

nursing diagnosis for activity intolerance: Manual of Nursing Diagnosis Marjory Gordon, 2014-11-13 Manual of Nursing Diagnosis, Thirteenth Edition outlines all the major and most recent diagnoses listed by the North American Nursing Diagnosis Association (NANDA) in order to help nurses create care plans for patients with any range of ailments. Updated and organized to meet the needs of both novices and expert diagnosticians, it includes sample admission assessment guides for the family, community, and individual. Manual of Nursing Diagnosis, Thirteenth Edition is an easy to use quick reference that shows nurses how to use diagnoses to guide questions and observations beyond the basic assessment. In addition, it helps nurses use diagnostic categories in other clinical activities, including critical paths and quality improvement. New to this edition are the latest NANDA changes, including: 21 new nursing diagnoses 9 revised nursing diagnoses 6 retired nursing diagnoses. --Publisher description.

nursing diagnosis for activity intolerance: Nursing Diagnosis Manual Marilyn E Doenges, Mary Frances Moorhouse, Alice C Murr, 2016-01-14 Here's the 5th Edition of the resource you'll turn to again and again to select the appropriate diagnosis and to plan, individualize, and document care for more than 850 diseases and disorders. A new, streamlined design makes reference easier than ever. Only in the Nursing Diagnosis Manual will you find for each diagnosis...defining characteristics presented subjectively and objectively - sample clinical

applications to ensure you have selected the appropriate diagnoses - prioritized action/interventions with rationales - a documentation section, and much more!

nursing diagnosis for activity intolerance: Nursing Care Plans & Documentation Lynda Juall Carpenito-Moyet, 2009 The Fifth Edition of Nursing Care Plans and Documentation provides nurses with a comprehensive guide to creating care plans and effectively documenting care. This user-friendly resource presents the most likely diagnoses and collaborative problems with step-by-step guidance on nursing action, and rationales for interventions. New chapters cover moral distress in nursing, improving hospitalized patient outcomes, and nursing diagnosis risk for compromised human dignity. The book includes over 70 care plans that translate theory into clinical practice. Online Tutoring powered by Smarthinking--Free online tutoring, powered by Smarthinking, gives students access to expert nursing and allied health science educators whose mission, like yours, is to achieve success. Students can access live tutoring support, critiques of written work, and other valuable tools.

nursing diagnosis for activity intolerance: *Handbook of Nursing Diagnosis* Lynda Juall Carpenito-Moyet, 2006 The newly revised Eleventh Edition of this best-selling handbook is an easy-to-carry, accessible guide to the latest NANDA-approved nursing diagnoses. Section 1 contains all nursing diagnoses, including definitions, characteristics, related factors, outcomes, and interventions. Section 2 contains Diagnostic Clusters with a collaborative focus. Features include Author's Notes, key concepts, interventions with rationales, focus assessment criteria, and outcome criteria. This edition includes listings of associated NIC (Nursing Interventions Classifications) and NOC (Nursing Outcomes Classifications) for every NANDA diagnosis. New diagnoses added and modified in accordance with the latest NANDA meeting are in an appendix for easy access.

nursing diagnosis for activity intolerance: Nursing Diagnosis Handbook Betty J. Ackley, Gail B. Ladwig, 2006 This bestselling reference helps users make nursing diagnoses and write care plans with ease and confidence. It helps nurses correlate nursing diagnoses with known information about clients on the basis of assessment findings, established medical or psychiatric diagnoses, and the current treatment plan. This extensively revised and updated edition presents the most up-to-date information on all 2005-2006 NANDA-approved nursing diagnoses. You'll also appreciate the handbook's integration of both the NIC and NOC taxonomies; evidence-based nursing interventions; home care; client/family teaching; and multicultural, geriatric, and pediatric considerations. Instructor resources are available; contact your sales representative for details. Helps users understand how to make a nursing diagnosis and plan care using the nursing process and evidence-based nursing research. The practical, step-by-step format helps users make diagnoses and write care plans with ease and confidence. Integrates both NIC and NOC interventions and outcomes in each care plan. Features essential home care, client/family teaching, multicultural, pediatric, and geriatric considerations relevant to today's nursing students. Helps users easily create customized plans of care using the Ackley, Ladwig Care Plan Constructor on the Evolve website. This edition has been extensively revised and updated to include the most up-to-date information on all 2005-2006 NANDA-approved nursing diagnoses. The new alphabetical thumb tabs offer easy access to specific nursing diagnosis care plans. Features a helpful explanation of how to use the Guide to Nursing Diagnoses and Guide to Planning Care sections. Features many enhancements to the accompanying Evolve website (see Ancillaries for details). Includes more detailed information about NOC and its use in nursing care planning. Guide to Nursing Diagnoses section features more emotional/spiritual elements and now includes coverage of Mad Cow, E. Coli, West Nile, Foodborne Illnesses, and Pesticide Contamination. Guide to Planning Care section now offers updated and expanded multicultural, pediatric, and evidence-based interventions/rationales. An appendix now includes Gordon's Functional Health Patterns, as adapted by the authors.

nursing diagnosis for activity intolerance: **Sparks and Taylor's Nursing Diagnosis Reference Manual** Sheila S. Ralph, Cynthia M. Taylor, 2013-01-03 Sparks and Taylor's Nursing Diagnosis Reference Manual 9th edition provides clearly written, authoritative care plan guidelines for all 2012-2014 NANDA International (NANDA-I) approved nursing diagnoses. The book is full of

bright colors, and organized by life-stages and type of care. Each Part opens with a new feature, Applying Evidence-Based Practice, which responds directly to the content. Each NANDA diagnosis includes associated Nursing Interventions Classifications (NIC) and Nursing Outcomes Classifications (NOC), and the nursing process is integrated throughout. This book is the ideal resource for any clinical setting.

nursing diagnosis for activity intolerance: Nursing Diagnosis & Intervention Gertrude K. McFarland, Elizabeth A. McFarlane, 1997 This book provides thorough coverage of both theory and practice of nursing diagnosis. It uses a narrative rather than a list format to explain nursing diagnosis. The book details the formulation of a nursing diagnosis and writing a care plan, as well as providing a resource to clear assessment parameters and planning care.

nursing diagnosis for activity intolerance: Maternal Newborn Nursing Care Plans Carol J. Green, 2016 Explains how to assess, plan, provide, and evaluate care for pregnancy, delivery, recovery, abnormal conditions, and newborn care. Features more than 65 of the most common and high-risk care plans for nursing care using the nursing process approach, this text includes NIC interventions, discussions on collaborative problems, key nursing activities, signs and symptoms, and diagnostic studies. --From publisher description.

nursing diagnosis for activity intolerance: Nursing Diagnosis Reference Manual Sheila Sparks Ralph, Cynthia M. Taylor, 2005 Nursing Diagnosis Reference Manual, Sixth Edition helps nursing students and practicing nurses prepare care plans accurately and efficiently for every NANDA-approved nursing diagnosis. The book features a life-cycle format, with sections on adult, adolescent, child, maternal-neonatal, and geriatric health. Sections on community-based health (care plans on home health, health promotion, and more) and psychiatric/mental health round out the volume. Each care plan includes clear-cut criteria for identifying the right nursing diagnosis, assessment guidelines, outcome statements, rationales with all interventions, and documentation guidelines.

nursing diagnosis for activity intolerance: All-in-One Care Planning Resource Pamela L. Swearingen, 2012-01-01 The only book featuring nursing care plans for all core clinical areas, Swearingen's All-In-One Nursing Care Planning Resource, 4th Edition provides 100 care plans with the nursing diagnoses and interventions you need to know to care for patients in all settings. It includes care plans for medical-surgical, maternity/OB, pediatrics, and psychiatric-mental health, so you can use just one book throughout your entire nursing curriculum. This edition includes a new care plan addressing normal labor and birth, a new full-color design, new QSEN safety icons, new quick-reference color tabs, and updates reflecting the latest NANDA-I nursing diagnoses and collaborative problems. Edited by nursing expert Pamela L. Swearingen, this book is known for its clear approach, easy-to-use format, and straightforward rationales. NANDA-I nursing diagnoses are incorporated throughout the text to keep you current with NANDA-I terminology and the latest diagnoses. Color-coded sections for medical-surgical, maternity, pediatric, and psychiatric-mental health nursing care plans make it easier to find information quickly. A consistent format for each care plan allows faster lookup of topics, with headings for Overview/Pathophysiology, Health Care Setting, Assessment, Diagnostic Tests, Nursing Diagnoses, Desired Outcomes, Interventions with Rationales, and Patient-Family Teaching and Discharge Planning. Prioritized nursing diagnoses are listed in order of importance and physiologic patient needs. A two-column format for nursing assessments/interventions and rationales makes it easier to scan information. Detailed rationales for each nursing intervention help you to apply concepts to specific patient situations in clinical practice. Outcome criteria with specific timelines help you to set realistic goals for nursing outcomes and provide quality, cost-effective care. NEW! Care plan for normal labor and birth addresses nursing care for the client experiencing normal labor and delivery. UPDATED content is written by practicing clinicians and covers the latest clinical developments, new pharmacologic treatments, patient safety considerations, and evidence-based practice guidelines. NEW full-color design makes the text more user friendly, and includes NEW color-coded tabs and improved cross-referencing and navigation aids for faster lookup of information. NEW! Leaf icon highlights coverage of

complementary and alternative therapies including information on over-the-counter herbal and other therapies and how these can interact with conventional medications.

nursing diagnosis for activity intolerance: All-In-One Care Planning Resource - E-Book Pamela L. Swearingen, 2015-02-02 NEW! Care plan for normal labor and birth addresses nursing care for the client experiencing normal labor and delivery. UPDATED content is written by practicing clinicians and covers the latest clinical developments, new pharmacologic treatments, patient safety considerations, and evidence-based practice guidelines. NEW full-color design makes the text more user friendly, and includes NEW color-coded tabs and improved cross-referencing and navigation aids for faster lookup of information. NEW! Leaf icon highlights coverage of complementary and alternative therapies including information on over-the-counter herbal and other therapies and how these can interact with conventional medications.

nursing diagnosis for activity intolerance: **Handbook of Medical-surgical Nursing** , 2006 Now in its updated Fourth Edition, this best-selling handbook is the most comprehensive pocket-sized guide to medical-surgical nursing. More than 300 disorders and treatments are covered in alphabetical order in a consistent quick-reference format, with icons, sidebars, alerts, illustrations, and tables to highlight key points. This edition covers bioterrorism-related disorders and new treatments such as enhanced external counterpulsation therapy and cardiac resynchronization therapy. A new icon highlights gender differences. Laboratory test results now include SI values. A new appendix on emergency preparedness is included.

nursing diagnosis for activity intolerance: *Pediatric Home Care for Nurses* Wendy Votroubek, 2010-09-15 Pediatric Home Care is a practice-based text perfect for either students or for supporting pediatric nurses practicing in a home-care setting. The text includes a variety of nursing information required for this type of care across a large spectrum of physiologic categories and acuity levels. The Third Edition has been completely revised and updated to reflect the most current practice and technology and includes a new focus on evidence based practice.

nursing diagnosis for activity intolerance: Alexander's Care of the Patient in Surgery - E-Book Jane C. Rothrock, 2014-03-01 NEW and UNIQUE! Patient-Centered Care boxes feature simplified conversations that you can have with patients to help them better understand their surgical procedure. NEW! A Critical Thinking Question at the end of every chapter tests your understanding of chapter content. NEW! Key Points at the end of every chapter help you retain important concepts from the text. NEW! Laboratory Values appendix contains all laboratory value information in one convenient location.

nursing diagnosis for activity intolerance: **Conceptual Care Mapping - E-Book** Barbara L. Yoost, Lynne R Crawford, 2017-01-08 Learn to prioritize patient care quickly and effectively! Conceptual Care Mapping: Case Studies for Improving Communication, Collaboration, and Care uses a 'two books in one' approach to guide you through the most important aspects of patient care and critical thinking. The first section explores the title's three C's of care, communication, and collaboration within the healthcare team. The second section includes case studies that progress from simple clinical conditions to the more complex. Conceptual Care Mapping provides the tools you need to understand and implement effective patient care plans. With mapping practice both online and in the text, this unique book is the ideal resource for any first-year nursing student! - Unique! Conceptual Care Mapping guides your care — using templates and an online, interactive conceptual care map creator — from the author team who developed this new method of organizing care information. - Unique! 30 case studies focus on the patients you are most likely to see in clinical experience as a first-year student — and are organized from simple to complex, guiding you through prioritization and teaching you to think critically. - Unique! Elsevier's Conceptual Care Map (CCM) creator helps you create, update, manage, and submit care maps quickly and effectively. - Unique! Answers and completed care maps are provided for selected cases to reinforce what you've learned. - Unique! Improving Patient Care section not only provides an overview of care mapping, but also focuses on three key elements of patient care: care, communication (both with other healthcare professionals and with patients), and collaboration. - Unique! Interprofessional Education (IPE) and

collaboration are key elements in all of the case studies, helping you understand how you will work within the larger healthcare team. - Full-color design makes it easier to match up your online care mapping with content in the text.

nursing diagnosis for activity intolerance: Maternity and Pediatric Nursing Susan Scott Ricci, Terri Kyle, 2009 Authors Susan Ricci and Terri Kyle have teamed up to deliver a unique resource for your students to understand the health needs of women and children. This new combination book, *Maternity and Pediatric Nursing*, will empower the reader to guide women and their children toward higher levels of wellness throughout the life cycle. The textbook emphasizes how to anticipate, identify, and address common problems to allow timely, evidence-based interventions. Features include unfolding case studies throughout each chapter, multiple examples of critical thinking, and an outstanding visual presentation with extensive illustrations depicting key concepts. A bound-in CD-ROM and a companion Website include video clips and NCLEX®-style review questions.

nursing diagnosis for activity intolerance: Clinical Aromatherapy E-Book Jane Buckle, 2003-08-20 Aromatherapy is one of the main complementary therapies to be practiced by nurses and other health care professionals in hospital, hospice, and community settings. Written by a nurse, this clinical text highlights how aromatherapy can enhance care and the role health care professionals play in its practice. It examines key facts and issues in aromatherapy practice, and applies these within a variety of contexts and conditions, taking a carefully holistic approach in dealing with the patient. - An introduction to the principles and practice of aromatherapy, including contraindications and toxicity. - Contains an in-depth clinical section dealing with the management of common problems such as infection and pain. - Gives examples of which specific oils might be used in treatment. - Illustrates the application of aromatherapy in specific clinical specialties, particularly nursing. - Draws on over 700 references - Includes a new chapter on psychiatric nursing and aromatherapy to include: depression, psychosis, bipolar, compulsive addictive, addiction and withdrawal.

Related to nursing diagnosis for activity intolerance

Nursing and midwifery - World Health Organization (WHO) Nursing encompasses autonomous and collaborative care of individuals of all ages, families, groups and communities, sick or well and in all settings. It includes the promotion of

Nursing and midwifery - World Health Organization (WHO) Nursing and midwifery occupations represent a significant share of the female workforce. More than 80% of the world's nurses work in countries that are home to half of the

State of the world's nursing report 2025 The 2025 edition of the State of the world's nursing provides the most comprehensive and up-to-date analysis of the nursing workforce. The report features new

Nursing workforce grows, but inequities threaten global health goals The global nursing workforce has grown from 27.9 million in 2018 to 29.8 million in 2023, but wide disparities in the availability of nurses remain across regions and countries,

State of the world's nursing 2020: investing in education, jobs and The State of the world's nursing 2020 report provides the latest, most up-to-date evidence on and policy options for the global nursing workforce

Countries advance "Nursing Action" initiative to tackle nurse The retention of nurses in Member States of the WHO European Region is central to "Nursing Action", a landmark European Union (EU)-funded initiative aimed at strengthening

State of the world's nursing 2025 - World Health Organization (WHO) State of the world's nursing 2025 provides updated data and evidence on the global nursing workforce. The level of data reflects a 33% increase in the number of countries reporting on a

Webinar - State of the World's Nursing 2025 (SoWN 2025) report WHO is currently developing the State of the World's Nursing 2025 (SoWN 2025) report, which will be launched on 12

Nursing EURO - World Health Organization (WHO) To safeguard the future health workforce and the provision of high-quality health care, steps must be taken to ensure that nursing and midwifery are seen as attractive career

Nursing and midwifery - World Health Organization (WHO) Nursing encompasses autonomous and collaborative care of individuals of all ages, families, groups and communities, sick or well and in all settings. It includes the promotion of

State of the world's nursing report 2025 The 2025 edition of the State of the world's nursing provides the most comprehensive and up-to-date analysis of the nursing workforce. The report features new

State of the world's nursing 2020: investing in education, jobs and The State of the world's nursing 2020 report provides the latest, most up-to-date evidence on and policy options for the global nursing workforce

Countries advance “Nursing Action” initiative to tackle nurse The retention of nurses in Member States of the WHO European Region is central to “Nursing Action”, a landmark European Union (EU)-funded initiative aimed at strengthening

State of the world's nursing 2025 - World Health Organization (WHO) State of the world's nursing 2025 provides updated data and evidence on the global nursing workforce. The level of data reflects a 33% increase in the number of countries reporting on a

Webinar - State of the World's Nursing 2025 (SoWN 2025) report WHO is currently developing the State of the World's Nursing 2025 (SoWN 2025) report, which will be launched on 12 May 2025. This report will offer an updated,

Nursing EURO - World Health Organization (WHO) To safeguard the future health workforce and the provision of high-quality health care, steps must be taken to ensure that nursing and midwifery are seen as attractive career

Nursing and midwifery - World Health Organization (WHO) Nursing encompasses autonomous and collaborative care of individuals of all ages, families, groups and communities, sick or well and in all settings. It includes the promotion of

Nursing and midwifery - World Health Organization (WHO) Nursing and midwifery occupations represent a significant share of the female workforce. More than 80% of the world's nurses work in countries that are home to half of the

State of the world's nursing report 2025 The 2025 edition of the State of the world's nursing provides the most comprehensive and up-to-date analysis of the nursing workforce. The report features new

Nursing workforce grows, but inequities threaten global health goals The global nursing workforce has grown from 27.9 million in 2018 to 29.8 million in 2023, but wide disparities in the availability of nurses remain across regions and countries,

State of the world's nursing 2020: investing in education, jobs and The State of the world's nursing 2020 report provides the latest, most up-to-date evidence on and policy options for the global nursing workforce

Countries advance "Nursing Action" initiative to tackle nurse

State of the world's nursing report 2025 The 2025 edition of the State of the world's nursing provides the most comprehensive and up-to-date analysis of the nursing workforce. The report features new

Nursing workforce grows, but inequities threaten global health goals The global nursing workforce has grown from 27.9 million in 2018 to 29.8 million in 2023, but wide disparities in the availability of nurses remain across regions and countries,

State of the world's nursing 2020: investing in education, jobs and The State of the world's nursing 2020 report provides the latest, most up-to-date evidence on and policy options for the global nursing workforce

Countries advance "Nursing Action" initiative to tackle nurse The retention of nurses in Member States of the WHO European Region is central to "Nursing Action", a landmark European Union (EU)-funded initiative aimed at strengthening

State of the world's nursing 2025 - World Health Organization State of the world's nursing 2025 provides updated data and evidence on the global nursing workforce. The level of data reflects a 33% increase in the number of countries reporting on a

Webinar - State of the World's Nursing 2025 (SoWN 2025) report WHO is currently developing the State of the World's Nursing 2025 (SoWN 2025) report, which will be launched on 12 May 2025. This report will offer an updated,

Nursing EURO - World Health Organization (WHO) To safeguard the future health workforce and the provision of high-quality health care, steps must be taken to ensure that nursing and midwifery are seen as attractive career

2025 2018 2790 2023 2980

Related Articles

- [sat practice test 1 pdf](#)
- [diagram of a sailing ship](#)
- [hayward pool pump diagram](#)

[Back to Home](#)